



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

testing of building stones, of the museums and collections of minerals and rocks within the limits of the kingdom, and of articles in the geology of Saxony.

W. S. B.

Notes. — The lavas of the early Tertiary volcanoes¹ of the Absaroka range on the east side of the Yellowstone National Park consist of a repeated succession of hornblendic and micaceous andesites, basalts, pyroxene, andesites, and finally a series of great flows of basalts. With these are associated immense deposits of tuffs, agglomerates, and igneous conglomerates.

Cushing² describes the augite-syenite gneiss near Loon Lake in the Adirondack district, New York, as medium grained, grayish green rocks composed of feldspar, pyroxene or hornblende, quartz, and sometimes biotite or garnet. They are undoubtedly metamorphosed intrusive rocks that are intimately associated with gneisses of sedimentary origin. The feldspar is usually a microperthite, but there are usually present in all slides small quantities of oligoclase. The pyroxene is principally augite, but hypersthene is often associated with it. The quartz is in elongated cylindrical individuals. The rocks are autoclastic in structure and are also foliated. In composition they are close to akerite.

SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	FeO	CaO	BaO	MgO	K ₂ O	Na ₂ O	P ₂ O ₅	Loss	Tot.
63.45	.07	18.31	.42	3.56	2.93	.13	.35	5.15	5.06	tr.	30	= 99.73

It is plain from the analysis that the augite is essentially a calcium, ferrous, aluminous variety unusually rich in alumina. The microperthite is approximately Or₃ AC₅. The syenite is thought to be closely related in origin to the anorthosite of the district.

Baalow³ gives excellent descriptions of some of the handsome autoclastic conglomerates met with in the Grenville and Hastings series of Ontario. They resemble very closely true conglomerates, but their genesis from banded rocks by dynamic agencies is clearly traced. The illustrations accompanying the descriptions are particularly interesting.

Pirssow⁴ collected together in a few pages the evidence that points to the conclusions that the phenocrysts of intrusive rocks are often formed in place, and are not intratelluric. The reasons for this con-

¹ Presidential Address of Arnold Hague, *Geol. Soc. of Washington*, 1899.

² *Bull. Geol. Soc. of America*, vol. x, p. 177.

³ *Ottawa Naturalist*, vol. xii, p. 205, 1899.

⁴ *Amer. Jour. Sci.*, vol. vii, p. 271, 1899.

clusion are briefly as follows: porphyritic intrusive masses often possess peripheral portions completely devoid of phenocrysts; dykes connected with large intrusions may be free from phenocrysts, while the mass of the intrusions is filled with them; flat crystals are often arranged haphazard in sheet and dykes and not in obedience to any law of flowage; phenocrysts often enclose crystals identical with those composing the matrix which surrounds them, and, finally, phenocrysts are often surrounded by microlites orientated parallel to the bounding faces of the large crystal, indicating that the latter was growing after the former had crystallized.

The conditions governing the consolidation and crystallization of igneous rocks are decrease in temperature, chemical composition of the magma, the influence of mineralizing vapors, pressure and increasing viscosity. In the view of the author "the greatest determinant in the formation of rock structure is the ratio of time in the fall of temperature between the point where the insolubility and crystallizing moment of a compound begins to the increasing viscosity."

The Inwood limestone¹ in the northern part of Manhattan Island is cut by pegmatite dykes, some of which are well exposed a few blocks north of Fort George. Near the contact the limestone contains tremolite, biotite, and brown tourmaline, while the last-named mineral occurs also in the peripheral portion of the pegmatite.

¹ Eckel. *Amer. Geologist*, vol. xxiii, p. 122, 1899.